

Contents

Preface	vii	
Abbreviations	xv	
1 Introduction to Plant Biochemistry	1	
2 Approaches to Understanding Metabolic Pathways	5	
What we need to understand a metabolic pathway	5	
Chromatography	7	
Electrophoresis	11	
The use of isotopes	14	
Current research techniques use a range of molecular biology approaches	16	
Unique aspects of plant metabolism and their impact on metabolic flux	26	
Metabolic control analysis theory	27	
Coarse and fine metabolic control	29	
Compartmentation: keeping competitive reactions apart	33	
Understanding plant metabolism in the individual cell	33	
The isolation of organelles	34	
Summary	36	
Further Reading	36	
3 Plant Cell Structure	39	
Cell structure is defined by membranes	40	
The plasma membrane	46	
Vacuoles and the tonoplast membrane	48	
The endomembrane system	49	
Cell walls serve to limit osmotic swelling of the enclosed protoplast	54	
The nucleus contains the cell's chromatin within a highly specialized structure, the nuclear envelope	56	
Mitochondria are ubiquitous organelles, which are the site of cellular respiration		57
Peroxisomes house vital biochemical pathways for many plant cell processes		58
Plastids		59
Summary		63
Further Reading		64
4 Light Reactions of Photosynthesis		65
Bacteria evolved the basic photochemical pathways found in plants today		65
Chlorophyll captures light energy and converts it to a flow of electrons		71
Carotenoids extend the spectral range of light that can be utilized in photosynthesis		74
Photosystem II splits water to form protons and oxygen, and reduces plastoquinone		75
The Q cycle uses plastoquinol to reduce plastocyanin and transport protons into the lumen		78
Photosystem I takes electrons from plastocyanin and reduces ferredoxin, which is used to make NADPH and other reduced compounds		80
ATP synthase utilizes the proton motive force to generate ATP		83
Cyclic photophosphorylation generates ATP independently of water oxidation and NADPH formation		85
Regulation of electron flow pathways in response to fluctuating light levels		86
Scavenging and removal of superoxides, peroxides, and other radicals by dismutases and antioxidants		87
Mechanisms for safely returning the levels of trapped high energy states to the ground state		88
Nonphotochemical quenching and the xanthophyll cycle		89
Summary		90
Further Reading		91

5 Photosynthetic Carbon Assimilation 93

Photosynthetic carbon assimilation produces most of the biomass on Earth	93
Carbon dioxide enters the leaf through stomata but water is also lost in the process	94
Carbon dioxide is converted to carbohydrates using energy derived from sunlight	94
The Calvin cycle is used by all photosynthetic eukaryotes to convert carbon dioxide to carbohydrate	96
Discovery of the Calvin cycle	96
There are three phases to the Calvin cycle	97
Calvin cycle intermediates may be used to make other photosynthetic products	108
The Calvin cycle is autocatalytic and produces more substrate than it consumes	108
Calvin cycle activity and regulation	109
Rubisco is a highly regulated enzyme	111
Rubisco oxygenase: the starting point for the photorespiratory pathway	113
The photorespiratory pathway: enzymes in the chloroplast, peroxisome, and mitochondria	113
The isolation and analysis of mutants and the photorespiratory pathway	117
Photorespiration may provide essential amino acids and protect against environmental stress	117
Photorespiration and the loss of photosynthetically fixed carbon	118
Photorespiration uses ATP and reductant	119
Decreasing global carbon dioxide concentrations caused a rapid evolution of C ₄ photosynthesis	119
C ₄ photosynthesis concentrates carbon dioxide at the active site of Rubisco	120
Spacial separation of the two carboxylases in C ₄ leaves	120
Stages of C ₄ photosynthesis and variations to the basic pathway	122
C ₃ –C ₄ intermediate species may represent an evolutionary stage between C ₃ and C ₄ plants	126
The C ₄ pathway can exist in single cells of some species	128
Some of the C ₄ pathway enzymes are light-regulated	129
Crassulacean acid metabolism as a feature of desert plants	130
Temporal separation of the carboxylases in CAM	130
Crassulacean acid metabolism as a flexible pathway	130

Phosphoenolpyruvate carboxylase in crassulacean acid metabolism plants regulated by protein phosphorylation	133
Crassulacean acid metabolism is thought to have evolved independently on several occasions	133
C ₃ , C ₄ , and CAM photosynthetic pathways: advantage and disadvantages	134
C ₃ , C ₄ , and CAM plants differ in their facility to discriminate between different isotopes of carbon	138
Summary	140
Further Reading	141

6 Respiration 143

Overview of respiration	143
The main components of plant respiration	144
Plants need energy and precursors for subsequent biosynthesis	144
Glycolysis is the major pathway that fuels respiration	145
Hexose sugars enter into glycolysis and are converted into fructose 1,6-bisphosphate	148
Fructose 1,6-bisphosphate is converted to pyruvate	148
Alternative reactions provide flexibility to plant glycolysis	149
Plant glycolysis is regulated by a bottom-up process	151
Glycolysis supplies energy and reducing power for biosynthetic reactions	151
The availability of oxygen determines the fate of pyruvate	151
The oxidative pentose phosphate pathway is an alternative catabolic route for glucose metabolism	153
The irreversible oxidative decarboxylation of glucose 6-phosphate generates NADPH	153
The second stage of the oxidative pentose phosphate pathway returns any excess pentose phosphates to glycolysis	153
All or part of the OPPP is duplicated in the plastids and cytosol	155
The tricarboxylic acid cycle is located in the mitochondria	155
Pyruvate oxidation marks the link between glycolysis and the tricarboxylic acid cycle	155
The product of pyruvate oxidation, acetyl CoA, enters the tricarboxylic acid cycle via the citrate synthase reaction	164
Substrates for the tricarboxylic acid cycle are derived mainly from carbohydrates	167
The tricarboxylic acid cycle serves a biosynthetic function in plants	167

Anaplerotic reactions are needed to enable intermediates to be withdrawn from the tricarboxylic acid cycle	169	7 Synthesis and Mobilization of Storage and Structural Carbohydrates	195
The tricarboxylic acid cycle is regulated at several steps	170	Role of carbohydrate metabolism in higher plants	195
Recent research into a thioredoxin/NADPH redox system for regulating tricarboxylic acid cycle enzymes and other mitochondrial proteins	172	Sucrose is the major form of carbohydrate transported from source to sink tissue	197
The mitochondrial electron transport chain oxidizes reducing equivalents produced in respiratory substrate oxidation and produces ATP	172	Sucrose phosphate synthase is an important control point in the sucrose biosynthetic pathway in plants	198
Main protein complexes of the electron transport chain	173	Sensing, signaling, and regulation of carbon metabolism by fructose 2,6-bisphosphate	200
Plant mitochondria possess additional respiratory proteins that provide a branched electron transport chain	175	Fructose 2,6-bisphosphate enables the cell to regulate the operation of multiple pathways of plant carbohydrate metabolism	200
Plant mitochondria contain four additional NAD(P)H dehydrogenases	176	Fructose 2,6-bisphosphate as a regulatory link between the chloroplast and the cytosol	204
The physiologic function of the alternative NAD(P)H dehydrogenases remains the subject of some speculation	176	Sucrose breakdown occurs via sucrose synthase and invertase	205
Plant mitochondria contain an alternative oxidase that transfers electrons from QH ₂ to oxygen and provides a bypass of the cytochrome oxidase branch	177	Starch is the principal storage carbohydrate in plants	209
The alternative oxidase is a dimer of two identical polypeptides with a nonheme iron center	178	Starch synthesis occurs in plastids of both source and sink tissues	209
Alternative oxidase isoforms in plants are encoded by discrete gene families	178	Starch formation occurs in water-insoluble starch granules in the plastids	213
Alternative oxidase activity regulated by 2-oxo acids and by reduction and oxidation	180	The composition and structure of starch affects the properties and functions of starches	215
The alternative oxidase adds flexibility to the operation of the mitochondrial electron transport chain	181	Starch degradation is different in different plant organs	216
The alternative oxidase may prevent the formation of damaging reactive oxygen species within the mitochondria	182	The nature and regulation of starch degradation is poorly understood	216
Alternative oxidase appears to play a role in the response of plants to environmental stresses	182	Transitory starch is remobilized initially by a starch modifying process that takes place at the granule surface during the dark period	218
Alternative oxidase and NADH oxidation	183	The regulation of starch degradation is unclear	219
Plant mitochondria and uncoupling proteins	183	Fructans are probably the most abundant storage carbohydrates in plants after starch and sucrose	220
ATP synthesis in plant mitochondria is coupled to the proton electrochemical gradient that forms during electron transport	183	A model has been proposed for the biosynthesis of the different fructan molecules found in plants	220
ATP synthase uses the proton motive force to generate ATP	184	Fructan-accumulating plants are abundant in temperate climate zones with seasonal drought or frost	222
Mitochondrial respiration interacts with photosynthesis and photorespiration in the light	187	Trehalose biosynthesis is not just limited to resurrection plants	222
Emerging research area into supercomplexes and metabolons	191	Trehalose synthesis in higher plants and its role in the regulation of carbon metabolism	223
Summary	191	Plant cell wall polysaccharides	224
Further Reading	192	Synthesis of cell wall sugars and polysaccharides	225
		Cellulose	225
		Matrix components consist of branched polysaccharides	228

Expansins and extensins, proteins that play both enzymatic and structural roles in cell expansion	234	The GS genes and proteins show discrete cellular localization and different responses to light and nutrients	260
Lignin	234	Glutamine synthetase activity is regulated by metabolites and effectors, and may be modified by phosphorylation and 14-3-3 binding	261
Summary	235	Further evidence of the functions of glutamine synthetase isoenzymes has come from studies of mutants and transgenic plants	261
Further Reading	235	Higher plants contain two forms of GOGAT, one is ferredoxin-dependent and the other is NADH-dependent	263
8 Nitrogen and Sulfur Metabolism	237	Both Fd- and NADH-GOGAT are located in the plastid and exist as monomeric proteins in most species	263
Nitrogen and sulfur must be assimilated in the plant	237	The tissue and cellular localization of Fd- and NADH-GOGAT provides a clue to their function in higher plants	264
Apart from oxygen, carbon, and hydrogen, nitrogen is the most abundant element in plants	238	Further evidence of the function of Fd- and NADH-GOGAT has come from the analysis of mutants and transgenic plants	264
Nitrogen fixation: some plants obtain nitrogen from the atmosphere via a symbiotic association with bacteria	239	Sulfur is an essential macronutrient but it represents only 0.1% of plant dry matter	265
Symbiotic nitrogen fixation involves a complex interaction between host plant and microorganism	242	Sulfate is relatively abundant in the environment and serves as a primary sulfur source for plants	266
Nodule-forming bacteria (<i>Rhizobiaceae</i>) are composed of the three genera <i>Rhizobium</i> , <i>Bradyrhizobium</i> , and <i>Azorhizobium</i>	242	The assimilation of sulfate	267
The nodule environment is generated by interaction between legume plant host and rhizobia	244	Adenosine 5'-phosphosulfate reductase is composed of two distinct domains	268
Nitrogen fixation is energy expensive, consuming up to 20% of total photosynthates generated	245	Sulfite reductase is similar in structure to nitrite reductase	269
Mycorrhizae are associations between soil fungi and plant roots that can enhance the nitrogen nutrition of the plant	246	Sulfation is an alternative minor assimilation pathway incorporating sulfate into organic compounds	269
Most higher plants obtain nitrogen from the soil in the form of nitrate	248	Amino acids synthesis is also essential for plant growth and development	270
In higher plants there are multiple nitrate carriers with distinct properties and regulation	249	Carbon flow is essential to maintain amino acid production	270
Nitrate reductase catalyzes the reduction of nitrate to nitrite in the cytosol of root and shoot cells	250	Depending on the plant species and tissue, nitrogen movement through the plant varies	272
The production of nitrite is rigidly controlled by the expression, catalytic activity, and degradation of NR	251	Aminotransferase reactions are central to amino acid metabolism by distributing nitrogen from glutamate to other amino acids	273
Nitrite reductase, localized in the plastids, catalyzes the reduction of nitrite to ammonium	253	Asparagine, aspartate, and alanine synthesis	275
Plant cells have the capacity for the transport of ammonium ions	255	Glycine and serine synthesis	276
Ammonium is assimilated into amino acids	258	The aspartate family of amino acids: lysine, threonine, isoleucine, and methionine	276
Ammonium originates from both primary and secondary sources	258	The branched chain amino acids valine and leucine	279
Ammonium is assimilated by glutamine synthetase and glutamate synthase, which combine together in the glutamine synthetase/ glutamate synthase cycle	259	Sulfur-containing amino acids cysteine and methionine	280
GS is an octameric protein with two isoforms, localized in the cytosol and plastid	259		

Glutamine, arginine, and proline synthesis	282	The products of the oxidation of lipids and the resulting metabolites are collectively known as oxylipins	320
The synthesis of the aromatic amino acids: phenylalanine, tyrosine, and tryptophan	284	The waxy cuticle coats all land plants	322
Histidine synthesis	285	Role of suberin as a hydrophobic layer	324
Large amounts of nitrogen can be present in nonprotein amino acids	285	Storage lipids are primarily a storage form of carbon and chemical energy	325
Plant storage proteins: why do plants store proteins and what sort of proteins do they store?	286	Release of fatty acids from acyl lipids	328
Vicilins and legumins are the main storage proteins in many dicotyledonous plants	288	The breakdown of fatty acids occurs via oxidation at the β carbon and subsequent removal of two carbon units	329
Prolamins are major storage proteins in cereals and grasses	290	Summary	333
2S albumins are important but minor components of seed proteins	292	Further Reading	334
Where are seed proteins synthesized and how do they reach their storage compartment?	292	10 Alkaloids	335
Protein stores are degraded and mobilized during seed germination	296	Plants produce a vast array of chemicals that deter or attract other organisms	335
Vegetative organs store proteins, which are very different from seed proteins	297	Alkaloids, a chemically diverse group that all contain nitrogen along with a number of carbon rings	336
Despite their diversity, storage proteins share common characteristics	299	Functions of alkaloids in plants and animals	336
Summary	299	The challenges and complexity of alkaloid biosynthetic pathways	336
Further Reading	300	Amino acids as precursors in the biosynthesis of alkaloids	338
9 Lipid Biosynthesis	303	Terpenoid indole alkaloids are made from tryptamine and the terpenoid secologanin	338
Overview of lipids	303	Isoquinoline alkaloids are produced from tyrosine and include many valuable drugs such as morphine and codeine	344
Fatty acid biosynthesis occurs through the sequential addition of two carbon units	307	Tropane alkaloids and nicotine are found mainly in the Solanaceae	349
The condensation of nine two-carbon units is necessary for the assembly of an 18C fatty acid	307	Pyrollizidine alkaloids are found in four main families	354
The acyl group of malonyl CoA is linked to acetyl CoA and then transferred to acyl carrier protein via a malonyl CoA:ACP transacylase	312	Purine alkaloids as popular stimulants in beverages, and as poisons and feeding deterrents against herbivores	355
For the assembly of an 18C fatty acid from acetyl CoA using type II fatty acid synthase, 48 reactions are necessary and at least 12 different proteins involved	312	The diversity of alkaloids has arisen through evolution driven by herbivore pressure	356
Acyl-ACP utilization in the plastid	314	Summary	360
Regulation of fatty acid formation	314	Further Reading	361
Source of NADPH and ATP to support fatty acid synthesis	315	11 Phenolics	363
The two pathways of glycerolipid synthesis	315	Plant phenolic compounds are a diverse group with a common aromatic ring structure and a range of biological functions	363
Phosphatidic acid, produced in the plastids or endoplasmic reticulum, is a central intermediate in glycerolipid synthesis	316	The simple phenolics	364
Lipids function in signaling and defense	318	The more complex phenolics include the flavonoids, which have a characteristic three-membered A, B, C ring structure	367

Lignin is a complex polymer formed mainly from monolignol units	369	Summary	397
The tannins are phenolic polymers that form complexes with proteins	370	Further Reading	398
Most plant phenolics are synthesized from phenylpropanoids	370	12 Terpenoids	399
The shikimic acid pathway provides the aromatic amino acid, phenylalanine, from which the phenylpropanoids are all derived	371	Terpenoids are a diverse group of essential oils that are formed from the fusion of five-carbon isoprene units	399
The core phenylpropanoid pathway provides the basic phenylpropanoid units that are used to make most of the phenolic compounds in plants	375	Terpenoids serve a wide range of biological functions	402
Flavonoids are produced from chalcones, formed from the condensation of <i>p</i> -coumaryl CoA and malonyl CoA	379	The biosynthesis of terpenoids	411
Simple phenolics from the basic phenylpropanoid pathway are used in the biosynthesis of the hydrolyzable tannins	391	Subcellular compartmentation is important in the regulation of terpenoid biosynthesis	426
Lignin is formed from monolignol subunits in a complex series of reactions that are still being unraveled	392	Summary	428
		Further Reading	428
		Colour plate section appears between pages xxx and xxx	