

CONTENTS

<i>Detailed Contents</i>	xiii
Chapter 1: Statistical Thermodynamics of Biological Molecules	1
Chapter 2: Protein Structure	33
Chapter 3: Evolution of Protein Sequence & Structure	89
Chapter 4: Protein Stability	149
Chapter 5: Protein Folding	199
Chapter 6: Binding, Allostery, and Cooperativity	255
Chapter 7: Enzyme Kinetics	313
Appendix A: Calculation of the Excess Heat Capacity Curve of Protein Thermal Denaturation	379
Appendix B: Calculation of the Average Helicity from the Partition Function	381
Appendix C: Solution of Rate Equations for a Two-State System	383
<i>Index</i>	387

DETAILED CONTENTS

Chapter 1 Statistical Thermodynamics of Biological Molecules **1**

1.1	BIOLOGICAL MOLECULES HAVE SPECIAL CHEMICAL PROPERTIES	1
	Water Shapes the Properties of All Biological Structures	1
	The Structures of Biological Macromolecules are Key to their Function	2
	The Structures of Proteins that Exist Today have been Selected by Evolution	2
	Interactions in Biomolecules are Weak	3
	Communication Occurs through Interactions between Molecules or within Molecules	3
1.2	STATISTICAL THERMODYNAMICS RELATES MICROSCOPIC INTERACTIONS TO MACROSCOPIC PROPERTIES	4
	States with the Same Energy are Equally Probable	4
	States with Lower Energies have a Higher Probability	6
1.3	THE ENERGY OF AN ISOLATED SYSTEM IS CONSTANT; THE ENTROPY INCREASES TOWARD A MAXIMUM	8
	The Energy is a Measure of Motion and Interactions in a System	8
	The Entropy is a Measure of the Number of States	9
	The Entropy can be Explicitly Related to Probabilities	10
	The Second Law is a Statement about Probability	12
1.4	THE GIBBS ENERGY IS A MINIMUM AT EQUILIBRIUM	13
	The Enthalpy is a Thermodynamic Function More Useful than the Energy in the Laboratory	13
	The Partition Function is Related to the Gibbs Energy	14
	The Gibbs Energy Provides the Criterion for Equilibrium at Constant T and P	16
1.5	THE GIBBS ENERGY DETERMINES THE CONCENTRATIONS AT EQUILIBRIUM	17
	The Chemical Potential of a Component is its Contribution to the Gibbs Energy of the System	17
	The Chemical Potential Increases with Concentration	18
	The Standard State must be Clearly Defined	19
	The Equilibrium Constant Provides a Measurement of the Standard Gibbs Energy Change	20
	Equilibrium Constants are Often Reported with Units (as a Reminder of the Standard State)	21
1.6	THE PARTITION FUNCTION IS A POWERFUL TOOL TO STUDY DISTRIBUTIONS OF MOLECULES	22
	Small Proteins are either Folded or Unfolded	23
	The Probabilities of Each State can be Obtained from the Partition Function	24
1.7	TWO-STATE SYSTEMS ARE COMMON IN CHEMISTRY	25
	Ribonuclease A is Almost Entirely Folded at Room Temperature	25
	The <i>cis</i> and <i>trans</i> Isomers of 2-butene are Populated to Different Extents at Equilibrium	27
	Nuclear Spins in a Magnetic Field have Almost Identical Populations	28
	The Partition Function Gives the Number of Occupied States	30

1.8	SUMMARY	30
1.9	PROBLEMS	31
1.10	FURTHER READING	32

Chapter 2 Protein Structure **33**

2.1	PROTEINS ARE POLYMERS OF A SPECIAL KIND	33
	Polypeptides are Formed by Amino Acids Connected by Peptide Bonds	33
	The Primary Structure of Proteins is the Sequence of Amino Acids	34
	Amino Acids are the Building Blocks of Proteins	34
	There are Twenty Amino Acids in Proteins	35
2.2	FLUORESCENCE SPECTROSCOPY IS A USEFUL TECHNIQUE TO MONITOR PROTEIN STRUCTURE AND INTERACTIONS	40
2.3	CHARGES IN A PROTEIN ARE DETERMINED BY THE IONIZATION OF ITS AMINO ACIDS	41
	The pK_a of a Chemical Group is a Measure of Its Proton (H^+) Binding Affinity	42
	One Ionizable Group has Two Forms, Depending on pH	42
	The Partition Function Allows a Rapid Calculation of the Fractions of Acidic and Basic Forms	44
	The Fractions of Protonated and Deprotonated States Change Sharply When the pH Approaches the pK_a	45
	Observable Properties of Molecular Ensembles are Weighted Averages of Individual Contributions	46
	The Problem with Two Ionizable Groups is Easy to Solve Using the Partition Function	47
2.4	THE SECONDARY STRUCTURE IS THE LOCAL SPATIAL ARRANGEMENT OF THE POLYPEPTIDE CHAIN	49
	The Peptide Group is Planar	49
	The Configuration About the Peptide Bond can be <i>cis</i> or <i>trans</i>	50
	Conformations are Changed by Rotations Around Single Bonds	50
	The Conformations of Residues in Peptides are Defined by the Torsion Angles ϕ and ψ	52
	The Ramachandran Diagram Shows the Values of ϕ and ψ that are Sterically Allowed	54
2.5	MOST OF THE SECONDARY STRUCTURE IN PROTEINS CONSISTS OF α -HELICES AND β -SHEETS	57
	The α -Helix is Stabilized by Hydrogen Bonds Between Residues i and $i + 4$.	57
	The β -Sheet is Stabilized by Hydrogen Bonds Between Adjacent Strands	59
	Circular Dichroism is a Simple Method to Identify α -Helices and β -Sheets	60
	Loops and Turns are Needed for Proteins to Fold to Compact Structures	61
	Globular Proteins Also Contain Another Type of Helix: the 3_{10} -Helix	62
	Collagen Adopts a Special Type of Helical Structure	63

6.7	LIGAND BINDING CHANGES PROTEIN STABILITY	304	7.7	SERINE PROTEASES USE ACID–BASE CATALYSIS TO HYDROLYZE PEPTIDE BONDS	344
	Specific Ligands Stabilize the Folded State	304		Substrate Specificity in Trypsin and Chymotrypsin is Conferred by the Residue Preceding the Bond Hydrolyzed	345
	Changes of pH may Stabilize Either the Folded or the Unfolded State	305		The Residues of the Catalytic Triad are the Most Important in the Mechanism of Serine Proteases	346
6.8	SUMMARY	307		The Catalytic Triad Accelerates Peptide Hydrolysis a Million Times	348
6.9	PROBLEMS	308			
6.10	FURTHER READING	310			
Chapter 7 Enzyme Kinetics		313			
7.1	ENZYMES ENHANCE REACTION RATES BY STABILIZING THE TRANSITION STATE	313	7.8	TRANSIENT KINETICS ALLOW DETERMINATION OF THE MOLECULAR RATE CONSTANTS	349
	Chemical Reactions Proceed through a Transition State, or Activated Complex	313		Binding Kinetics can be Determined Exactly in Favorable Circumstances	349
	Enzymes are Optimized to Bind the Transition State	315		If Binding is not Rapid Compared to the Chemical Step, Transient Kinetics Still Allow Determination of Rate Constants	351
	Enzymes Provide a Reaction Pathway through a Series of Stabilized Intermediates	317		Intermediates in the Mechanism of Enzyme-Catalyzed Reactions can be Detected in Transient Kinetics	354
7.2	LIGANDS WITH MULTIPLE CONTACTS REDUCE THE UNFAVORABLE ENTROPY LOSS UPON BINDING	318		The Serine Proteases Follow a Ping-Pong Mechanism with an Acyl-Enzyme Covalent Intermediate	355
	Many Ligands Interact with Proteins at Multiple Sites	318		“Burst” Kinetics Indicate that an Intermediate is Formed	356
	The Coupling Gibbs Energy Depends on the Number of Intervening Bonds	321			
7.3	THE DISTINCTIVE CONCEPT OF ENZYME KINETICS IS THE FORMATION OF AN ENZYME–SUBSTRATE COMPLEX	325	7.9	THE ACTIVITY OF ENZYMES DEPENDS ON PH	357
	Formation of a Michaelis–Menten Complex Leads to Saturation in the Reaction Kinetics	325		Enzymes are Often Synthesized as Inactive Precursors that are Post-Translationally Activated	357
	The Michaelis–Menten Equation Describes Steady-State Enzyme Kinetics	327		The pH Dependence of Enzyme Activity is Often Bell-Shaped	359
	K_M is the Concentration of Substrate at which the Rate is Half-maximum	328		The pH Determines How Much of the Active form of the Enzyme Exists	360
	The Catalytic Constant k_{cat} Measures the Rate at which the Enzyme Transforms the Bound Substrate	329		The Fraction of the Active form of Chymotrypsin and the Kinetic Parameters k_{cat} , K_M , and k_{cat}/K_M can be Calculated from the Partition Function	360
	The Specificity Constant k_{cat}/K_M Measures the Catalytic Efficiency of the Enzyme	330	7.10	CONFORMATIONAL DYNAMICS ARE ESSENTIAL IN THE MECHANISM OF DIHYDROFOLATE REDUCTASE	365
	The Constant k_{cat}/K_M Can Never Exceed the Diffusion Limit	331		Dihydrofolate Reductase Catalyzes a Hydride Transfer Reaction	365
	The Double-Reciprocal Lineweaver–Burk Plot is Useful in Qualitative Diagnostic of Enzyme Mechanisms and Inhibition	331		DHFR Cycles through Several Conformations During Catalysis	367
7.4	THE MICHAELIS–MENTEN EQUATION APPLIES TO MOST ENZYME REACTION MECHANISMS	333		The Rates of All DHFR Reactions were Determined by Transient Kinetics Experiments	368
	The K_M and k_{cat} are in General Combinations of Rate Constants and Concentrations	333		A Kinetic Isotope Effect Occurs when a C–H Bond is Broken in the Rate-Limiting Step	370
	The Rates of Complex Mechanisms can be Determined by the King–Altman Method	333		Protein Conformational Dynamics Determine the Rates of Hydride Transfer and Ligand Release	371
7.5	MOST ENZYME REACTIONS INVOLVE TWO SUBSTRATES	336	7.11	SUMMARY	372
	In Sequential Reactions all Substrates Bind Before Products are Formed	336	7.12	PROBLEMS	373
	In Ping-Pong Reactions Substrate Binding Alternates with Product Release	337	7.13	FURTHER READING	376
7.6	THE PARTITION FUNCTION CAN BE USED TO OBTAIN STEADY-STATE KINETICS IN SPECIAL CASES	339			
	The Michaelis Constant Behaves as an Apparent Dissociation Constant	339			
	Enzyme Inhibition Kinetics can be Derived from the Partition Function	340			
				Appendix A Calculation of the Excess Heat Capacity Curve of Protein Thermal Denaturation	379
				Appendix B Calculation of the Average Helicity from the Partition Function	381
				Appendix C Solution of Rate Equations for a Two-State System	383