

Contents in Detail

Preface	vii		
Acknowledgments	viii		
1 Levels of Gene Control	1		
Introduction	1		
1.1 Protein content varies among different cell types	1		
Specific methods can be used to study the expression of individual proteins in tissues and cells	1		
General methods can be used for studying the overall protein composition of tissues and cells	2		
1.2 The mRNA content varies among different cell types	6		
Specific methods can be used to study the expression of individual mRNAs in different tissues and cells	6		
General methods can be used to study the overall population of mRNAs expressed in different tissues and cells	7		
1.3 The DNA content of different cell types is generally the same	9		
Specific methods can be used to study individual genes in different tissues and cells	10		
General methods can be used to study the total DNA in different tissues and cells	10		
Exceptional cases do exist in which changes to the DNA occur in specific tissues or cell types	13		
1.4 Transcriptional or post-transcriptional control?	15		
Studies of nuclear RNA suggest that gene transcription is regulated	16		
Pulse labeling studies directly demonstrate transcriptional control	18		
Nuclear run-on assays allow transcriptional control to be demonstrated for a wide range of genes	19		
Polytene chromosomes provide further evidence for transcriptional control	21		
Transcriptional control can operate at the level of chromatin structure and at the level of production of the primary RNA transcript	22		
1.5 Small RNAs and the regulation of gene expression	23		
miRNAs are processed from a single-stranded precursor which folds to form a double-stranded hairpin loop	24		
Many siRNAs are processed from a double-stranded precursor	25		
Conclusions	27		
Key concepts	27		
Further reading	28		
2 Structure of Chromatin	29		
Introduction	29		
Regulation of transcription in eukaryotes is much more complex than in prokaryotes	29		
2.1 Commitment to the differentiated state and its stability	30		
Cells can remain committed to a particular differentiated state even in the absence of its phenotypic characteristics	30		
Cells can become committed to a particular differentiated state prior to actual phenotypic differentiation	31		
2.2 The nucleosome	33		
The nucleosome is the basic unit of chromatin structure	33		
Nucleosome structure or position can be altered by chromatin-remodeling processes	36		
2.3 Histone modifications and histone variants	37		
Histones are subject to a variety of post-translational modifications	37		
Histone variants are encoded by distinct genes to those encoding the standard histone isoforms	43		
2.4 The 30 nm chromatin fiber	44		
The 30 nm fiber represents a further compaction of the beads-on-a-string structure	44		
Histone H1 and post-translational modifications of the other histones are involved in the formation of the 30 nm fiber	45		

2.5 Structural and functional domains in chromatin	47	3.5 Changes in chromatin structure in the regulatory regions of active or potentially active genes	78
The 30 nm fiber is further compacted by looping	47	DNaseI-hypersensitive sites can be identified in active or potentially active genes	78
Locus-control regions regulate the chromatin structure of a large region of DNA	48	DNaseI-hypersensitive sites frequently correspond to regulatory DNA sequences	80
Insulators block the inappropriate spread of particular chromatin structures	50	DNaseI-hypersensitive sites represent areas which are either nucleosome-free or have an altered nucleosomal structure	81
Heterochromatin is a very tightly packed form of chromatin	51	Chromatin remodeling can be produced by proteins capable of displacing nucleosomes or altering their structure	82
The chromosome is the visible result of chromatin compaction	52	The SWI-SNF and NURF chromatin-remodeling complexes are recruited to the DNA by a variety of different mechanisms	84
Conclusions	52	3.6 Other situations in which chromatin structure is regulated	85
Key concepts	53	In female mammals one of the two X chromosomes is inactivated	86
Further reading	54	The active and inactive X chromosomes have a different chromatin structure	86
		The XIST regulatory RNA is specifically transcribed on the inactive X chromosome	87
3 Role of Chromatin Structure in Gene Control	55	Genomic imprinting involves the specific inactivation of either the maternally or paternally inherited copy of specific genes	88
Introduction	55	Imprinting involves changes in chromatin structure	90
3.1 Changes in chromatin structure in active or potentially active genes	56	Conclusions	92
Active DNA is organized in a nucleosomal structure	56	Key concepts	93
Active or potentially active chromatin shows enhanced sensitivity to DNaseI digestion	57	Further reading	93
3.2 Alterations in DNA methylation in active or potentially active genes	59	4 The Process of Transcription	95
Decreased DNA methylation is associated with active or potentially active genes	59	Introduction	95
DNA methylation plays a key role in regulating chromatin structure	62	4.1 Transcription by RNA polymerases	95
DNA methylation patterns can be propagated stably through cell divisions	62	Transcription by RNA polymerase I is relatively simple	96
DNA methylation recruits inhibitory proteins that produce a tightly packed chromatin structure	64	Transcription by RNA polymerase III is more complex than for RNA polymerase I	96
3.3 Modification of histones in the chromatin of active or potentially active genes	66	Transcription by RNA polymerase II is much more complex than transcription by RNA polymerases I and III	99
Acetylation	66	Transcription by the three different polymerases has a number of common features	101
Methylation	68	Transcription takes place in defined regions of the nucleus	105
Ubiquitination and sumoylation	72	4.2 Transcriptional elongation and termination	107
Phosphorylation	73	Transcriptional elongation requires further phosphorylation of RNA polymerase II	107
3.4 Interaction of different histone modifications, DNA methylation, and RNAi	74		
The different histone modifications interact functionally with one another	74		
Histone modifications interact with DNA methylation to regulate chromatin structure	75		
RNAi can induce alterations in chromatin structure	76		

Termination of transcription occurs downstream of the polyadenylation signal	109	The leucine zipper is a dimerization domain which allows DNA binding by the adjacent basic domain	147
4.3 The gene promoter	110	In some transcription factors, the basic DNA-binding domain is found associated with a helix-loop-helix dimerization domain	148
The 70 kDa heat-shock protein gene contains a typical promoter for RNA polymerase II	111	Dimerization between factors provides an additional level of regulation	149
The hsp70 gene promoter contains several DNA sequence motifs which are found in a variety of other gene promoters	112	Other domains can also mediate DNA binding	150
The heat-shock element is found only in heat-inducible genes	112	5.2 Activation of transcription	152
Other response elements are found in the promoters of genes with different patterns of expression	114	Activation domains can be identified by “domain-swap” experiments	152
The proteins binding to short DNA sequence elements can be characterized by a variety of techniques	116	Several different classes of activation domain exist	154
Promoter regulatory elements act by binding factors which either affect chromatin structure and/or influence transcription directly	120	How is transcription activated?	155
4.4 Enhancers and silencers	121	Activators can interact with TFIID	156
Enhancers are regulatory sequences that act at a distance to increase gene expression	121	Activators can interact with TFIIB	157
Many enhancers have cell-type- or tissue-specific activity	123	Activators can interact with the mediator and SAGA complexes	157
Proteins bound at enhancers can interact with promoter-bound factors and/or alter chromatin structure	125	Activators can interact with co-activators	158
Silencers can act at a distance to inhibit gene expression	128	Activators can interact with modulators of chromatin structure	160
Conclusions	130	Activators have a multitude of targets	161
Key concepts	131	5.3 Repression of transcription	162
Further reading	132	Repressors can act indirectly by inhibiting the positive effect of activators	162
5 Transcription Factors and Transcriptional Control	133	Repressors can act directly by inhibiting the assembly or activity of the basal transcriptional complex	164
Introduction	133	5.4 Regulation at transcriptional elongation	166
5.1 DNA binding by transcription factors	135	Regulation of transcription can occur at the elongation stage, as well as at initiation	166
The helix-turn-helix motif is found in a number of transcription factors which regulate gene expression during embryonic development	136	Factors which regulate transcriptional elongation target the C-terminal domain of RNA polymerase II	168
The helix-turn-helix domain found in homeodomain proteins is a DNA-binding domain	137	5.5 Regulation of transcription by RNA polymerases I and III	169
In the POU domain transcription factors, the homeodomain forms part of a larger DNA-binding motif	139	Transcription by RNA polymerases I and III can be regulated by alterations in chromatin structure	170
The two-cysteine–two-histidine (Cys ₂ His ₂) zinc finger is found in multiple copies in many transcription factors	141	Transcription by RNA polymerases I and III can be regulated by altering the expression or activity of components of the basal transcriptional complex	170
The nuclear receptors contain two copies of a multi-cysteine zinc finger distinct from the Cys ₂ His ₂ zinc finger	143	Regulation of transcription by RNA polymerase III can involve specific transcription factors binding to RNA as well as to DNA	171
		Conclusions	172
		Key concepts	172
		Further reading	173

6 Post-transcriptional Processes	175		
Introduction	175		
6.1 Capping	175		
The capping process modifies the 5' end of the RNA transcript	175		
The cap enhances translation of the mRNA by the ribosome	176		
6.2 Polyadenylation	178		
The polyadenylation process modifies the 3' end of the RNA transcript	178		
Polyadenylation enhances the stability of the mRNA	178		
6.3 RNA splicing	180		
RNA splicing removes intervening sequences and joins exons together	180		
Specific RNAs and proteins catalyze the process of RNA splicing	180		
6.4 Coupling of transcription and RNA processing within the nucleus	185		
Transcriptional initiation and elongation are coupled to post-transcriptional processes	185		
Post-transcriptional processes can interact with one another	186		
6.5 RNA transport	187		
RNA transport is coupled to other post-transcriptional processes	187		
6.6 Translation	189		
Translation of the mRNA takes place on cytoplasmic ribosomes	189		
Translational initiation involves initiation factors binding to the cap	190		
Translational elongation involves base-pairing of triplet codons in the mRNA with tRNA anticodons	191		
Translational termination occurs at specific stop codons	194		
6.7 RNA degradation	195		
RNA degradation occurs in both the nucleus and the cytoplasm	195		
RNA degradation in the cytoplasm involves prior de-adenylation and decapping of the mRNA	196		
Conclusions	198		
Key concepts	198		
Further reading	199		
7 Post-transcriptional Regulation	201		
Introduction	201		
7.1 Alternative RNA splicing	202		
RNA splicing can be regulated	202		
		Alternative splicing represents a major regulatory process which supplements transcriptional control	202
		Alternative RNA splicing involves specific splicing factors that promote or inhibit the use of specific splice sites	209
		Factors regulating alternative splicing have been identified by genetic and biochemical methods	210
		The processes of transcription and alternative splicing interact with one another	214
		Alternative RNA splicing is a very widely used method of supplementing transcriptional control	215
7.2 RNA editing	217		
		Some cases of RNA editing involve a change from a C to a U residue	217
		Other cases of RNA editing involve a change from an A to an I residue	217
7.3 Regulation of RNA transport	219		
		Specific proteins can regulate the transport of individual mRNAs from nucleus to cytoplasm	219
		RNA transport processes can also regulate the location of individual mRNAs within the cytoplasm	221
7.4 Regulation of RNA stability	223		
		Gene regulation can involve alterations in RNA stability	223
		Specific sequences in the mRNA are involved in the regulation of its stability	223
		RNA stability changes supplement transcriptional control in cases where a rapid response is required	225
7.5 Regulation of translation	226		
		Translational control occurs in specific situations such as fertilization	226
		Translational control can involve either modifications in the cellular translational apparatus or specific proteins which recognize sequences in the target RNA	226
		Translational control can be produced by modifications in the cellular translation apparatus	227
		Translational control can be produced by proteins binding to specific sequences in the RNA itself	230
		Translational control frequently occurs when a rapid response is required but also occurs for the genes encoding some transcription factors	235
7.6 Post-transcriptional inhibition of gene expression by small RNAs	235		
		Small RNAs can inhibit gene expression post-transcriptionally	235

Small RNAs can induce mRNA degradation	236	Transcription factors can be activated by cleavage of a membrane-bound precursor	260
Small RNAs can repress mRNA translation	237	Cleavage of a transcription factor can convert it from an activator to a repressor	262
miRNAs regulate gene expression at multiple levels	240	Cleavage of a lipid link can be used to activate a transcription factor	262
Conclusions	241	8.5 Regulation of post-transcriptional processes by cellular signaling pathways	263
Key concepts	241	The PI3-kinase/Akt system plays a key role in regulating gene expression in response to growth factors or insulin	263
Further reading	242	Akt regulates RNA splicing by phosphorylating splicing factors	264
8 Gene Control and Cellular Signaling Pathways	243	Akt regulates mRNA translation via the TOR kinase, which phosphorylates proteins involved in translation	264
Introduction	243	Akt/TOR can also stimulate mRNA translation by enhancing the transcription of genes encoding RNAs and proteins involved in protein synthesis	265
Transcription factors can be regulated by controlling their synthesis or by controlling their activity	243	A variety of kinases inhibit translation by phosphorylating eIF2	266
Multiple mechanisms regulate transcription factor activity	245	Individual kinases can produce multi-level regulation of gene expression	267
8.1 Regulation of transcription factor activity by ligands which enter the cell	246	Conclusions	268
Transcription factors can be activated by direct binding of ligands which enter the cell	246	Key concepts	271
Members of the nuclear receptor family of transcription factors are activated by binding of the appropriate ligand	246	Further reading	272
Following ligand-mediated activation, the glucocorticoid receptor can repress as well as activate gene transcription	249	9 Gene Control in Embryonic Development	273
The HSF is activated by stressful stimuli and induces the transcription of genes encoding protective proteins	251	Introduction	273
8.2 Regulation of transcription factor activity by phosphorylation induced by extracellular signaling molecules	252	Regulation of mRNA translation occurs following fertilization	273
Transcription factors can be phosphorylated by receptor-associated kinases	252	Transcriptional control processes activate the embryonic genome	274
Transcription factors can be phosphorylated by kinases activated by specific intracellular second messengers such as cyclic AMP	252	The Oct4 and Cdx2 transcription factors regulate the differentiation of ICM and trophectoderm cells	276
Transcription factors can be phosphorylated by signaling cascades consisting of several protein kinases	254	9.1 Regulation of gene expression in pluripotent ES cells	277
Transcription factor activity can be regulated by phosphorylation of an inhibitory protein: the NF κ B/I κ B system	255	ES cells can differentiate into a wide variety of cell types	277
8.3 Regulation of transcription factor activity by other post-translational modifications	257	Several transcription factors are specifically expressed in ES cells and together can reprogram differentiated cells to an ES-cell-like phenotype	279
Acetylation	257	ES-cell-specific transcription factors can activate or repress the expression of their target genes	279
Methylation	258	ES-cell-specific transcription factors regulate genes encoding chromatin-modifying enzymes and miRNAs	281
Ubiquitination and sumoylation	259	The REST transcription factor plays a key role in down-regulating the expression of	
8.4 Regulation of transcription factor activity by signals which regulate precursor processing	260		
Transcription factors can be activated by cleavage of a precursor which contains an inhibitory region	260		

ES-cell-specific transcription factors during differentiation	282	MyoD is a basic helix-loop-helix transcription factor which is able to regulate gene expression	306
ES cells have an unusual pattern of histone methylation	282	MyoD is regulated by controlling both its synthesis and its activity	307
The polycomb complex regulates histone methylation in ES cells	284	Other muscle-specific transcription factors can induce muscle cell differentiation	308
Polycomb protein complexes regulate the expression of miRNA genes in ES cells	286	MEF2 is a downstream regulator of muscle-cell specific gene transcription	310
Chromatin structure in ES cells is regulated by multiple effects on histones	286	10.2 Regulation of gene expression in neuronal cells	313
9.2 Role of gene regulation in the development of <i>Drosophila melanogaster</i>	288	Basic helix-loop-helix transcription factors are also involved in neuronal differentiation	313
A gradient in expression of the Bicoid transcription factor defines the anterior–posterior axis in the early <i>Drosophila</i> embryo	288	The REST transcription factor represses the expression of neuronal genes	315
Bicoid activates a cascade of genes encoding other transcription factors, producing a segmented pattern of <i>Eve</i> gene expression	289	Neuronal cells express specific alternative splicing factors	317
The Bicoid system involves both transcriptional and post-transcriptional regulation	290	Translational control plays a key role in synaptic plasticity in neuronal cells	319
Homeodomain transcription factors specify segment identity in the <i>Drosophila</i> embryo	291	miRNAs play a key role in the regulation of neuronal gene expression	320
Protein–protein interactions control the effect of homeodomain-containing transcription factors on gene expression	292	10.3 Regulation of yeast mating type	322
9.3 Role of homeodomain factors in mammalian development	292	Yeast cells can be a or α in mating type	322
Homeodomain transcription factors are also found in mammals	292	Mating-type switching is controlled by regulating the transcription of the <i>HO</i> gene	323
Mammalian <i>Hox</i> genes are expressed in specific regions of the developing embryo	293	The SBF transcription factor activates <i>HO</i> transcription only in the G1 phase of the cell cycle	323
Transcription of individual <i>Hox</i> genes is regulated by gene-specific regulatory regions	294	The <i>Ash-1</i> transcription factor represses <i>HO</i> transcription in daughter cells	324
<i>Hox</i> gene transcription is also dependent on the position of the gene in the <i>Hox</i> gene cluster	295	The a and α gene products are homeodomain-containing transcription factors	324
<i>Hox</i> gene expression is also regulated at the post-transcriptional level by specific miRNAs	296	The α1 and α2 proteins interact with the MCM1 transcription factor to respectively activate α -specific genes and repress a -specific genes	325
Differential regulation of different <i>Hox</i> genes by Sonic Hedgehog controls the differentiation of cells in the neural tube	296	The a1 factor plays a key role in repressing haploid-specific genes in diploid cells	326
Regulation of <i>Hox</i> gene expression by Sonic Hedgehog is also involved in limb formation	298	The yeast mating-type system offers insights of relevance to multicellular organisms	327
Conclusions	299	Conclusions	329
Key concepts	300	Key concepts	330
Further reading	301	Further reading	331
10 Control of Cell-type-specific Gene Expression	303	11 Gene Regulation and Cancer	333
Introduction	303	Introduction	333
10.1 Regulation of gene expression in skeletal muscle cells	305	11.1 Gene regulation and cancer	333
The MyoD protein can induce muscle cell differentiation	305	Oncogenes were originally identified in cancer-causing viruses	333
		Cellular proto-oncogenes are present in the genome of normal cells	334
		Cellular proto-oncogenes can cause cancer when they are over-expressed or mutated	336

Viruses can induce elevated expression of oncogenes	337	12.2 Chromatin structure and human disease	367
Proto-oncogene expression can also be enhanced by cellular mechanisms	338	DNA methylation	367
A variety of cellular mechanisms mediate enhanced expression of proto-oncogenes in different cancers	338	Histone-modifying enzymes	368
11.2 Transcription factors as oncogenes	340	Chromatin-remodeling complexes	369
The Fos and Jun oncogene proteins are cellular transcription factors which can cause cancer when over-expressed	340	12.3 Post-transcriptional processes and human disease	370
The <i>v-erbA</i> oncogene protein is a mutant form of the cellular thyroid hormone receptor	342	RNA splicing	370
Other transcription factor-related oncogenes are over-expressed due to chromosomal translocations	344	RNA translation	373
Chromosomal translocations can also produce novel oncogenic fusion proteins involving transcription factors	345	12.4 Infectious diseases and cellular gene expression	373
11.3 Anti-oncogenes	348	12.5 Gene regulation and therapy of human disease	375
Anti-oncogenes encode proteins which restrain cellular growth	348	Therapy could be achieved by altering the expression of transcription factors	375
The p53 protein is a DNA-binding transcription factor	349	Therapy could be achieved by altering the activity of transcription factors	376
The retinoblastoma protein interacts with other proteins to regulate transcription	352	Therapy could be achieved by targeting proteins which alter chromatin structure	378
Other anti-oncogene proteins also regulate transcription	354	Therapy could be achieved using designer zinc fingers to alter gene transcription	378
11.4 Regulation of gene expression: the relationship of cancer and normal cellular function	356	Therapy could be achieved by modulating RNA splicing	379
Oncogenes and anti-oncogenes interact to regulate the expression of genes encoding proteins which control cellular growth	356	Conclusions	381
Oncogenes and anti-oncogenes interact to regulate the expression of RNAs and proteins involved in mRNA translation	358	Key concepts	381
Oncogenes and anti-oncogenes interact to regulate the expression of microRNAs	359	Further reading	382
Conclusions	360	13 Conclusions and Future Prospects	383
Key concepts	361	Conclusions and future prospects	383
Further reading	362	Transcription factors interact with one another to regulate transcription	383
12 Gene Regulation and Human Disease	363	Transcription factors can repress gene expression as well as activating it	384
Introduction	363	DNA-binding transcription factors interact with co-activators/co-repressors and with regulators of chromatin structure	384
12.1 Transcription and human disease	363	Histone modifications play a central role in the regulation of chromatin structure	385
DNA-binding transcription factors	363	Co-activators/co-repressors link together different signaling pathways by interacting with multiple transcription factors	386
DNA-binding sites for specific transcription factors	364	Gene regulation is highly complex and involves both transcriptional and post-transcriptional regulation	386
Transcriptional co-activators	366	RNA molecules play a central role in regulating gene expression	387
Components of the basal transcriptional complex	367	Alterations in regulatory RNAs and proteins cause human disease	388
Factors involved in transcription by RNA polymerases I and III	367	Regulatory networks control gene expression	388
		Further reading	390
		Glossary	391
		Index	409